


Capsulolabroligamentous Structures of the Hip: Anatomy and Pathology

Michelle M. Nguyen, MD
VA Palo Alto Medical Center
Dept of Radiology, MSK Division

Lecture Outline

- I. Normal Articular Anatomy of the Hip
- II. Pathology of Joint Capsule and Ligaments
- III. Labrum:
 - I. Normal Anatomy
 - II. Pathology
 - III. Pitfalls
- IV. Miscellaneous
 - I. Supraacetabular fossa/Stellate lesion

Hip Joint: Articular Anatomy



Hip Joint

- Synovial ball-and-socket joint
- Rotate around 3 planes
 - Transverse axis: flexion and extension
 - Longitudinal axis: medial and lateral rotation
 - Sagittal axis: abduction and adduction



[http:// wikipedia.org](http://wikipedia.org)


Hip Joint: Osseous Anatomy

- **Femoral Head**

- Half a sphere
- Directed upward, medially and slightly anteriorly
- Smooth surface xc for roughened pit, the fovea

- **Femoral Neck**


- Entire anterior surface intracapsular
- Medial third of posterior surface intracapsular



Hip Joint: Osseous Anatomy

Acetabulum

- Cup-shaped socket
- Elevated bony rim with lunate cartilaginous surface
- Bony rim/cartilage absent inferiorly at the acetabular notch
- Acetabular fossa
 - Devoid of cartilage
 - Filled with fat
 - Covered by synovium




Hip Joint: Articular Anatomy

- **Labrum**

- Fibrocartilaginous rim attached to edge of the acetabulum
- Deepens the acetabular side of the joint
- Ends and merges inferiorly with the transverse ligament


- **Transverse Ligament**

- Spans the acetabular notch, converting notch to foramen
- A non-cartilaginous extension of the labrum





Ligamentum Teres Femoris

- Pyramidal ligament
- Covered by synovium
- Extends from femoral fovea as round band
- Inserts distally as 2 flattened bands on either side of acetabular notch, blending with fibers of transverse ligament
- Sometimes only synovial covering is present
- Contributes to stability in fetal and neonatal hips⁹
- Debated function in adult hips



Ligamentum Teres Femoris

- Pyramidal ligament
- Covered by synovium
- Extends from femoral fovea as round band
- Inserts distally as 2 flattened bands on either side of acetabular notch, blending with fibers of transverse ligament
- Sometimes only synovial covering is present
- Contributes to stability in fetal and neonatal hips⁹
- Debated function in adult hips



Ligamentum Teres Femoris

Table I. Proposed functions of the ligamentum teres (LT)


Function	Arguments for	Arguments against
Mechanical		
Stability	Rich in collagen ⁹ stiffness comparable to other ligaments, ⁹ hip joint stability gradually decreases during gestation, ²⁰ while the collagenous component of LT increases ¹⁰	Presence of numerous static and dynamic stabilisers around the hip; minimal increase in range of movement after sectioning the LT ⁷
Proprioception/co-ordination	Presence of FNE* in LT; ^{29,30} episodes of recurrent subluxation in athletes with injured LTs ²⁶ may be caused by loss of fine coordination	FNE identified in 100% of normal ²⁹ but only 66.6% of dysplastic ³⁰ hips; cause-and-effect relationship between subluxation and injury to LT not proven
Biological		
Nociception	Presence of FNE; ^{29,30} pain reported in otherwise healthy hips with ruptured LTs; ^{3,5} pain in inflammatory/degenerative arthropathy	Multiple potential sources of pain in inflammatory/degenerative arthropathy
Vascularity to femoral head	Vessels present in LT ^{10,65-67}	Penetration of fovea by vascular canals limited and extremely variable ¹⁰
Distribution of synovial fluid	None	Lack of experimental proof ¹³

* FNE, free nerve endings

Hip Joint: Synovial Cavity

Synovial Membrane


- Covers portion of femoral neck, femoral head articular surface, ligamentum teres
- Intimate with deep fibers of capsular ligaments
- Covers acetabular labrum and fossa



Hip Joint: Capsule

Fibrous capsule

- Circular and longitudinal fibers
- Strongest anteriorly and superiorly
- Attachments:
 - Proximal: acetabular margin (5-6mm above labrum), labrum, transverse ligament
 - Anterior/distal: surrounds femoral neck to trochanteric line
 - Posterior/distal: femoral neck to 1cm above trochanteric crest




Chatha DS et al. Mag Res Imag Clin N Am
2005;13: 605-15.

Hip Joint: Capsule

Zona orbicularis

- Deep circular fibers of capsule
- Collar about femoral neck
- No connection to bone
- Blends with the more superficial capsular ligaments
- Abundant at inferior and posterior capsule
- Tension at extreme flexion or extension
- Hip stability in distraction⁷
- “femoral arcuate ligament^{8,10”}



www.prohealthsys.com


Capsular Ligaments

- Hip joint capsule is intimate with and reinforced by capsular ligaments
- Strongest anteriorly

- Anterior Ligaments:

- Pubofemoral Ligament


- Iliofemoral Ligament



Anterior Capsular Ligaments

● Iliofemoral Ligament


- Ligament of Bigelow
- Strongest capsular ligament
- “Y” shaped
 - Apex attached to or near lower portion of anterior inferior iliac spine (AIIS)
 - Base attached along trochanteric line
- Two bands (majority)
 - Lateral band (L)
 - Medial band (M)
- Possible 3 bands⁸



Iliofemoral Ligament

- Main restrictor of external rotation and extension
- **Medial Band (A)**
 - Anatomy
 - Origin: btwn AIIIS and acetabular rim
 - Insertion: more distal trochanteric line of femoral neck
 - Almost vertical orientation
 - Function
 - Inhibitor of external rotation, especially in extension

*Martin H et al. Arthroscopy
2008; 24(2): 188-195*



Cadaveric analysis of anatomy and function:
Internal and external rotation thru ROM fr 30°
flexion to 10° extension along neutral swing path

Iliofemoral Ligament

Lateral Band (B)


● Anatomy

- Origin: AIIS
- Insertion: superior portion of trochanteric line
- More horizontal oblique orientation along femoral neck


● Function

- Main inhibitor of external rotation, along neutral swing
- Also inhibitor of internal rotation, esp in extension

*Martin H et al. Arthroscopy
2008; 24(2): 188-195*




Iliofemoral Ligament



Moorman CT et al. *J Bone Joint Surg Am.* 2003;85:1190-1196.


Anterior Capsular Ligaments: Pubofemoral Ligament

- Resembles a sling
- Anatomy¹²
 - Origin: Wide attachment to iliopectineal eminence, superior ramus and obturator crest
 - Insertion: Blends with capsule and deep surface of medial band of iliofemoral ligament. Attaches to femoral neck
- Function¹²
 - Inhibitor of external rotation in extension and abduction



Anterior Capsule: Inherent Weak Area

- Inherent weak area of anterior capsule between the medial band of the iliofemoral ligament and the pubofemoral ligament
- Potential communication between hip joint and iliopsoas bursa




Posterior Capsular Ischiofemoral Ligament

Anatomy¹²

- Origin: Ischial portion of acetabular rim and extends superolaterally as 2 bands
- Superior Band
 - Spirals across posterior femoral neck, blends with zona orbicularis
 - Inserts medial to anterosuperior base of greater trochanter
- Inferior Band
 - Crosses posterior femoral neck
 - Inserts medial to base of greater trochanter along posterior trochanteric crest

Function¹²: Inhibitor of external rotation in extension




Capsule: Normal Arthrogram

Anterior Joint Capsule


- Blue arrows
- Extends to trochanteric line

Posterior Joint Capsule

- Black arrowheads
- More medial and proximal than anterior joint capsule



Hip Joint: Normal MR Anatomy




Chatha DS et al. MR Imaging of Normal Hip. MRI Clin N Am 2005;13: 605-15.

Coronal T1FS Anterior to Posterior

Iliofemoral
ligament →

← Iliopsoas T.





Iliofemoral
ligament
(lateral)

Anterior
Labrum


Iliopsoas t.


Iliofemoral
ligament
(medial)

Iliofemoral
ligament

Transverse
ligament

Iliofemoral
ligament
(medial)






Labrum

Iliofemoral
ligament

Fovea

Transverse
ligament

Region of
pubofemoral
ligament



Labrum


Iliofemoral
ligament

Ligamentum
Teres

ZO

Transverse
ligament

ZO



Labrum

Iliofemoral
ligament

ZO


Acetabular
fossa

Ligamentum
Teres

Transverse
ligament

Synovial
longitudinal fibers/
retinacular
vessels

ZO



Labrum


Ischiofemoral
ligament

ZO

Ligamentum
Teres

Transverse
ligament

ZO




Labrum

Ischiofemoral
ligament

Labrum

ZO




ZO


Posterior
Labrum

Obturator
Externus
Tendon


Sagittal T1FS Medial to Lateral

Transverse
ligament






Transverse
ligament



Iliopsoas t.

Labrum


Labrum



Labrum

Iliopsoas t.

Labrum




Labrum

Iliofemoral
ligament

Labrum


Ischiofemoral
ligament



Iliofemoral
ligament

This is a grayscale MRI scan of a knee joint in a flexed position. The femur is at the top, and the tibia is at the bottom. The patella is visible in the center. Two ligaments are highlighted with white arrows and labels: the Iliofemoral ligament on the left and the Ischiofemoral ligament on the right. The background is dark, and the joint structures are in shades of gray.


Ischiofemoral
ligament



Iliofemoral
ligament

Ischiofemoral
ligament

ZO




Iliofemoral
ligament

Ischiofemoral
ligament

ZO

Obturator
Externus
Tendon




Iliofemoral
ligament

ZO


Obturator
Externus
Tendon

Axial T1FS Superior to Inferior




Labrum

Iliofemoral
ligament



Labrum

Iliofemoral
ligament



Iliofemoral
ligament


Labrum


Acetabular
Fossa

Fovea

Ischiofemoral
ligament

Labrum





Iliofemoral
ligament


Labrum

Acetabular
Fossa

Ligamentum
Teres

Ischiofemoral
ligament

Labrum




Iliofemoral
ligament

Ligamentum
Teres

ZO

Ischiofemoral
ligament

Labrum



Iliofemoral
ligament


Transverse
ligament

ZO

Ligamentum
Teres

ZO

Ischiofemoral
ligament




Iliofemoral
ligament

ZO

Transverse
ligament

ZO

Ischiofemoral
ligament



Ilio-femoral
ligament



Ischio-femoral
ligament



Pathology of Capsule and Ligaments




Capsular Laxity and Hip Instability

- Traumatic (acute): Dislocation or subluxation
- Atraumatic (non-acute):
 - Overuse (chronic repetitive microtrauma)
 - Connective tissue disorders (Marfan, Ehlers-Danlos)
- Capsular ligaments (iliofemoral ligament) and labrum are static stabilizers of hip
 - Once injured, the psoas major muscle (dynamic stabilizer) constantly contracts to provide hip stability
 - Leads to stiffness, flexion contractures of hip, back pain
- Treatment: Thermal capsulorrhaphy +/- plication and labral repair

Traumatic Posterior Subluxation

- Seen in athletes (American football)
- Fall on flexed adducted hip
- Low-energy trauma
- Difficult clinical diagnosis
- Avulsion fracture of posterior acetabulum
- Disruption of iliofemoral ligament and anterior capsule
- Hemarthrosis
- Risks: AVN, chondrolysis




Moorman CT et al. JBJS 2003;
85A(7):1190-6.

Traumatic Posterior Subluxation

THE JOURNAL OF BONE & JOINT SURGERY · JBJS.ORG
VOLUME 85-A · NUMBER 7 · JULY 2003


TRAUMATIC POSTERIOR HIP SUBLUXATION
IN AMERICAN FOOTBALL



Traumatic Posterior Subluxation

THE JOURNAL OF BONE & JOINT SURGERY · JBJS.ORG
VOLUME 85-A · NUMBER 7 · JULY 2003

TRAUMATIC POSTERIOR HIP SUBLUXATION
IN AMERICAN FOOTBALL



Patient #1



Patient #2



Hip Dislocation

- Posterior >> anterior dislocation
- Joint capsule is stronger anteriorly
- High-force impact
 - Flexed adducted internally rotated hip → Posterior dislocation
 - Abducted externally rotated hip → Anterior dislocation
- Risks: AVN, post-traumatic osteoarthritis




Philippon M et al. Arthroscopy
2009;25(2):169-174

Hip Dislocation


Imaging Findings

- Iliofemoral ligament disruption
 - 16 of 18 pts with traumatic dislocation¹⁴
- Labral and chondral injuries +/- entrapment
- Capsular/Ischiofemoral ligament disruption
- Ligamentum teres disruption¹⁵
- Femoral head or acetabular fracture
- Marrow and muscle edema
- Sciatic nerve injury

Ligamentous Injuries in Dislocation



Iliofemoral Ligament Disruption: contrast extends outside capsule



Ischiofemoral Ligament Tear: contrast extends outside posterior capsule



Traumatic Hip Dislocation in Professional Athletes¹⁵

TABLE 2. Hip Pathologies Identified at Arthroscopy

Patient No.	Months*	Spontaneous Reduction	Hip Pathology Identified at Arthroscopy						
			Labral	Adhesions	Capsule Tear	Ligamentum Teres	Cam	Pincer	Chondral Defect
1	7.3	N	Y	—	—	Y	Y	Y	Y
2	4.4	N	Y	—	Y	Y	Y	—	Y
3	4.0	N	Y	—	—	—	—	—	Y
4	3.3	N	Y	—	—	Y	—	—	Y
5	0.1	N	Y	—	—	Y	—	—	Y
6	3.1	N	Y	—	—	Y	—	—	Y
7	0	N	Y	—	—	Y	—	—	Y
8	6.6	Y	Y	—	—	—	—	Y	Y
9	1.5	N	Y	Y	Y	Y	Y	—	Y
10	2.1	N	Y	—	—	Y	Y	Y	Y
11	0.8	Y	Y	—	—	Y	Y	Y	Y
12	0.6	N	Y	—	—	Y	Y	—	Y
13	4.6	N	Y	Y	—	Y	Y	—	Y
14	18.5	N	Y	—	—	Y	Y	Y	Y

*Months from dislocation to hip arthroscopy.


Ligamentum Teres Injuries

Gray and Villar Classification¹⁷

- Type I Complete Rupture:
 - Usually acute and associated with major trauma/surgery
 - Often associated with other significant abnormalities (labral or chondral injuries)
- Type II Partial Rupture:
 - Usually associated with chronic hip pain
 - Often seen in athletes (ballet dancers, martial arts or high impact sports).
 - Third most common finding at arthroscopy in young athletes
- Type III Degenerative:
 - Usually associated with osteoarthritis in older patients

Traumatic Ligament Teres Tear: A source of Hip Pain¹⁶

- Mechanism⁹:
 - Fall on flexed adducted hip
 - Twisting injury, ext rotation
 - Hyperabduction
- Clinical Symptoms:
 - Deep anterior groin pain
 - Locking, clicking, popping
- 23 cases of traumatic tears of LT (15 major trauma, 8 twisting)
- 2 detected by preop MRA (7 MRA, 20 MRI done)
- Young pts (avg age=28)
- 8 cases were isolated findings.
- Treatment:
 - Debridement of torn fibers
 - Improved sx after treatmt






Byrd T et al. Arthroscopy 2004;20(4): 385-391

MR Imaging of Ligamentum Teres

MR Arthrogram^{18,19}:


- Axial oblique planes may be optimal
- Normal: Homogeneous low SI on both T1 and T2W
- Partial Tear: Intermed/high T2 signal and abnormal morphology
- Hypertrophied ligament: 2 mm medial extension beyond foveal insertion on axial oblique images.
 - If ligament is hypertrophied, either abnormal T2 signal or morphology determines partial tear.
- Occasionally will see bony intra-articular fragments from avulsion injuries of LT (more common with complete/type I injuries)

Ligamentum Teres Tears



24 yo professional soccer player with left hip pain


Coronal T2FS



Courtesy of Drs Craig Stewart and Tudor Hughes

24 yo professional soccer player with left hip pain

Axial T2 FS



24 yo professional soccer player with left hip pain


Axial T2 FS

Arthroscopic Findings:

Type 2 partial tear of left
ligamentum teres at its
foveal attachment



Labrum: Normal Anatomy and Pathology




Labrum

- Fibrocartilaginous structure:

- Attached to acetabular perimeter
- Inferiorly, attached to transverse acetabular ligament
- Collagen fibers contiguous with those in transverse ligament

- Morphology:


- Anterior labrum: Wider and thinner
- Posterior labrum: Less wide and thicker



my.statdx.com

Labrum



- **Function:**
 - Provides seal, keeping fluid btwn femoral head and acetabulum during mild compression
 - Helps contain femoral head during extreme ROM, esp hip flexion
- **Histology²:**
 - Dense connective tissue
 - Fibrocartilage
 - Collagen fibers (longitudinal circumferential bundles)
 - Neuroreceptors within labrum provide proprioception of hip⁴



Labrum: Vascularity

- Most vascular at peripheral 1/3 of labrum²
- Kelly et al³ divided labrum into capsular zone IA/IB and articular zone IIA/IIB
 - Capsular zone most vascular d/t abundant blood supply from joint capsule
 - Some blood supply from adj bone
 - Least blood supply in central and articular labrum

Peripheral labral tears near capsulolabral junction may heal better.



Kelly et al. Arthroscopy 2005;21(1):3-11


Shindle et al. JBJS 2008;90 Suppl 4:2-19

Labrum: MR Evaluation

- MR arthrography is sensitive and specific in detecting labral pathology
 - Debate that noncontrast MRI just as accurate³³
- Radial imaging has been suggested as better way to evaluate labrum.

Labrum Types: MR Imaging^{21,26}

- Triangular 69-80%
- Round 13-16%
- Irregular/Flat 7-13%
- “Absent” 1-2.5%
 - May not have been seen d/t technique
 - Degenerative or torn, not truly absent^{22,23}




Aydingoz U et al. Eur Radiol 2001;11:567-574.

Normal Labrum: MR Imaging

MR Findings:

- Triangular (most common)
- Homogeneous dark signal intensity on all sequences
- Closely attached to articular cartilage
- Clock face: sagittal plane²⁵
 - 12:00 superior
 - 3:00 anterior
 - 6:00 inferior
 - 9:00 posterior



Steinbach L et al. RadioGraphics
2002; 22:1223-1246

Normal Labrum: MR Imaging

MR Findings:

- Triangular (most common)
- Homogeneous dark signal intensity on all sequences
- Closely attached to articular cartilage
- Clock face: sagittal plane²⁵
 - 12:00 superior
 - 3:00 anterior
 - 6:00 inferior
 - 9:00 posterior



Labral Abnormalities/Tears

Causes

1. Degeneration
 - Intrasubstance abnl signal
2. Femoroacetabular Impingement
 - Cam type: femoral head/neck offset
 - Pincer type: abnormal acetabular morphology
 - Mixed type: most common
 - Developmental hip dysplasia
 - Coxa magna
3. Trauma
4. Repetitive activity in athletes


Clinical Findings:

- Asymptomatic: >70 yo
- Hip/Groin Pain
- Clicking

Resnick D. IDJ Advanced Intensive
MR Imaging Course Feb 2009

Femoroacetabular Impingement: Cam type

- Pain w/flexion and internal rotation and limited ROM w/ internal rotation
- Offset at femoral head/neck junction causes impingement on labrum with flexion
- “Pistol grip” deformity
- Young athletic persons
- Labral and chondral abnormalities
- Alpha angle $> 55^\circ$
- Herniation pits may be early osteoarthritis²⁹




Normal (a/b): Labrum clears with flexion

Cam type (c/d): Femoral head/neck offset impinges on acetabular labrum

Femoroacetabular Impingement: Pincer-type

- Abnormal acetabular morphology
 - Acetabular protrusion
 - Acetabular retroversion
 - Os acetabuli
 - Trauma
- Older female patients
- Over-coverage of femoral head → linear contact on acetabular labrum
- Mostly labral abnormalities
- Contrecoup chondral injury in posteroinferior acetabulum and femoral head



Normal (a/b): Labrum clears with flexion

Pincer type (e/f): Over-acetabular coverage of femoral head. Abnormal linear contact on acetabular labrum on flexion


Hong R, Hughes T, Gentili A, Chung C.
J of Mag Res Imag 2008;27:435

Acetabular Protrusion




Measure fr medial acetabular wall to ilioischial line

Acetabular Protrusio $> 3\text{mm}$ in men; $> 6\text{mm}$ in women



MR measurement B is the distance between the medial most point of the acetabular fossa and a line perpendicular to the horizontal axis that passes through the lateral most point of the posterior inner pelvic wall (curved arrow) at the level of the ischial spine.


Acetabular Retroversion



Anterior acetabular rim is more lateral and crosses over posterior acetabular rim. “Figure of 8” or crossover sign


Labral Tears Classification

- Arthroscopic classification by Lage²⁷
 - Radial flap
 - Radial fibrillated
 - Longitudinal peripheral
 - Unstable
- Initial MR Imaging classification by Czerny et al²⁸
 - I. Intralabral signal
 - II. Contrast extends into labrum
 - III. Labral detachment
- BUT, these 2 systems do not correlate well with ea other²⁵



Resnick's MR Classification of Labral Tears

- Use sagittal plane for location of tear
- Coronal plane (cross reference to other planes) for type of tear
- **Types:**
 1. Longitudinal vertical
 2. Longitudinal horizontal
 3. Radial
 4. Complex
 5. Labral detachment
 - From bone or articular cartilage
 6. Acetabulolabral articular destruction (ALAD)




Labral Tears: Location is Everything


- Most tears are anterior or anterosuperior
 - Theories: poorer vascular supply, mechanically weaker and subject to greater stress
 - Marginal attachment to cartilage, parallel collagen fibers to cartilage³²
- Posterosuperior tears can occur in younger pts and more common in Japan
- Normal labral variants are more likely in the lower quadrant

14 yo female ballerina with “snapping” sensation of right hip

MR Arthrogram: Coronal T1 FS





Axial T1 FS




Anterosuperior Labral Tear

68 yo woman with anterosuperior labral tear



38 yo male with avulsion of anterosuperior labrum



Dinauer et al. AJR 2004;183:1745-53

Labral Tears: Paralabral Cyst



38 yo female with right hip pain for 7 yrs, no trauma

Courtesy of Dr Tudor Hughes


Labral Tears: Paralabral Cyst



38 yo female with right hip pain for 7 yrs, no trauma

Courtesy of Dr Tudor Hughes

Acetabulolabral articular destruction (ALAD)




40 yo male with right hip pain

Courtesy of Dr Tudor Hughes

Labral Imaging Pitfalls

Terminology:

- Perilabral recess or sulcus: btwn labrum and capsule
 - NORMAL
- Sublabral recess or sulcus: btwn labrum and articular cartilage or subchondral bone
 - Normal or not??




Normal Perilabral Recess⁵

Sublabral Sulcus: Confusing Topic


- Breakdown of literature

- 1981 Walker³⁴: anterosuperior labral cleft in fetal labrum seen w/ light microscopy but not macroscopically, possible prep artifact?
- 1994 Keene³⁸: 100 pts arthroscopy. 2 posteroinferior sublabral sulci.
- 1996 Petersilge³⁵: 10 MRA. No sublabral sulcus.
- 1999 Czerny³⁶: 40 MRA; 6 cadavers. No sublabral sulcus.
- 2004 Dinauer³⁷: 54 MRA, 4 MRI. 13 Posteroinferior sublabral sulci.
- 2006 Saddik³⁹: 24 noncon MRI, 3 MRA. 12 anterosuperior, 14 posteroinferior, 2 anteroinferior, 2 posterosuperior sulci
- 2008 Studler⁴⁰: 57 MRA. 7 anteroinferior, 1 anterosuperior, 2 anterior sulci.
- Definition per above: Partial separation of labrum fr articular cartilage with normal underlying labrum. Gold standard was arthroscopy.
- BUT, there were some examples of sublabral sulcus on MR proven to be tear on arthroscopy

Posteroinferior Sublabral Sulcus³⁷




Posteroinferior Sublabral Sulcus³⁹




Posterosuperior Sublabral Sulcus³⁹




Anterosuperior Sublabral Sulcus³⁹




Anteroinferior Sublabral Sulcus⁴⁰



Tear or Sublabral Sulcus³⁹?



Tear or Sublabral Sulcus³⁹?




Anterosuperior
labral tear on
arthroscopy

Tear or Sublabral Sulcus³⁷?



Tear or Sublabral Sulcus³⁷?




Anterior labral
tear

Posterior
sublabral sulcus

Sublabral Sulcus: Bottom Line

- Describe the labral finding.
- Location is key.
- If in anterosuperior labrum, consider a tear.
- If in posteroinferior labrum, likely a sulcus since most labral tears are anterosuperior in location.
- In other quadrants, be careful dismissing finding as normal sulcus.

**Miscellaneous:
Supraacetabular Fossa
Stellate Crease/Lesion**



Supraacetabular Fossa and the Stellate Crease or Lesion

Supraacetabular Fossa

- Normal notch within the acetabulum deficient of cartilage
- Located above the acetabular fossa at 12:00
- Usually seen in young people

Stellate Crease

- Normal area of stellate appearance of deficient cartilage at or near supraacetabular fossa, seen on arthroscopy





FIG. 27. The stellate crease is frequently found directly superior to the acetabular fossa (AF) characterized by a stellate pattern of chondromalacia (arrows). This appears to be a normally occurring process, even in young adults, without clear prognostic significance. (Courtesy of J. W. Thomas Byrd, MD.)

Stellate Lesion


- A subchondral osseous fragment within the stellate crease
- A plica attached to the medial aspect of the fragment, extending medially and inferiorly
- May be symptomatic
- Hip plica described in arthroscopy literature⁴¹



Courtesy of Drs Resnick and Sampatchalit

Stoller D. MRI in Orthopaedics and Sports Medicine 3rd Edit.

Summary

- Normal anatomy and pathology of the capsule, ligaments and labrum of the hip
 - Pitfalls in imaging of the labrum
 - Stellate lesion
- 

References

1. Shindle M et al. Arthroscopic Management of Labral Tears in the Hip. J Bone Joint Surg Am. 2008;90 Suppl 4:2-19
2. Petersen W et al. Structure and Vascularity of the Labrum. Arch Orthop Trauma Surg 2003;123: 283–288
3. Kelly B et al. Vascularity of the Hip Labrum. Arthroscopy 2005; 21(1):3-11
4. Kim YT Azuma H. The nerve endings of the acetabular labrum. Clin Orthop Relat Res. 1995;320:176-81.
5. Chatha DS et al. MR Imaging of Normal Hip. MRI Clin N Am 2005;13: 605-15.
6. Meislin R et al. Role of Hip MRI in Management of Sports-related Injuries. MRI Clin N Am 2005;13: 635-40.
7. Ito et al. Proximal Hip Joint Capsule and Zona Orbicularis Contribute to Hip Joint Stability. J of Ortho Res 2009 on-line paper.
8. Fuss FK, Bacher A. New aspects of the morphology and function of the human hip joint ligaments. Am J Anat 1991;192:1-13.
9. Bardakos N et al. The Ligamentum Teres of the Adult Hip. J Bone Joint Surg 2009;91- B:8-15.
10. Hewitt J et al. Mech Properties of Human Hip Capsule Ligaments. J of Arthroplasty 2002;17(1): 82-9.
11. Moorman CT et al. Traumatic Posterior Hip Subluxation in American Football. J Bone Joint Surg Am. 2003;85:1190-1196.
12. Martin H et al. Function of Hip Capsular Ligaments: Quantitative Report. Arthroscopy 2008; 24(2): 188-195

References

13. Hong R, Hughes T, Gentili A, Chung C. Magnetic Resonance of the Hip. *J of Mag Res Imag* 2008;27:435-45.
14. Laorr A et al. Traumatic Hip Dislocation:Early MRI Findings. *Skel Rad* 1995;24:239
15. Philippon M et al. Arthroscopic Findings in Traumatic Hip Dislocation in 14 Professional Athletes. *Arthroscopy* 2009;25 (2):169-174.
16. Byrd T et al. Traumatic Rupture of Ligamentum Teres as a Source of Hip Pain. *Athroscopy* 2004;20(4): 385-391.
17. Gray AJR, Villar RN. The Ligamentum Teres of the Hip: An Arthroscopic Classification of Its Pathology. *Arthroscopy* 1997;13(5): 575-578.
18. Philippon MJ, Armfield DR, et al. Detection of Partial Tears of the Ligamentum Teres of the Hip Using MR Arthrography. Paper #346. AAOS Meeting, Thursday March 23, 2006
19. Armfield DR et al. Radiographic and MR Imaging of the Athletic Hip. *Clin of Sport Med* 2006(25):211-239
20. Shindle M et al. Diagnosis and Management of Traumatic and Atraumatic Hip Instability in the Athletic Patient. *Clin of Sport Med* 2006(25)-309-326
21. Aydingoz U, Ozturk MH. MR imaging of the acetabular labrum: a comparative study of both hips in 180 asymptomatic volunteers. *Eur Radiol* 2001;11:567-574.
22. Blankerbaker D et al. The Painful Hip: New Concepts. *Skeletal Radiol* 2006;35: 352-370
23. Dinauer P et al. Sublabral Sulcus at the Posteroinferior Acetabulum. *AJR* 2004;183:1745-1753

References

24. Steinbach L et al. MR Arthrography. *RadioGraphics* 2002; 22:1223–1246
25. Blackenbaker D et al. Classification and localization of acetabular labral tears. *Skel Rad* 2007; 36:391–397
26. Abe I et al. Acetabular Labrum: Abnormal Findings at MR Imaging in Asymptomatic Hips. *Radiology* 2000;216(2):576-81.
27. Lage L et al. The Acetabular Labral Tear: An Arthroscopic Classification. *Arthroscopy* 1996;12:269-72.
28. Czerny C et al. Lesions of Acetabular Labrum: Accuracy of MRI and MR Arthrography in Detecting and Staging *Radiology* 1996;200:225-30.
29. Leunig M et al. Fibrocystic changes at anterosuperior femoral neck: prevalence in hips with femoroacetabular impingement. *Radiology* 2005;236:237–246.
30. Tannast M et al. Femoroacetabular Impingement Radiographic Diagnosis—What the Radiologist Should Know. *AJR* 2007;188(6):1540-52.
31. Chen L et al. Acetabulum Protrusio and Center Edge Angle:New MR-Imaging Measurement Criteria. *Skel Rad* 2009(38):123-129.
32. Cashin M et al. Embryology of Acetabular Labral Chondral Complex. *JBJS* 2008 90-B(8):1019-24.
33. Mintz D et al. Magnetic Resonance Imaging of the Hip: Detection of Labral and Chondral Abnormalities Using Noncontrast Imaging. *Arthroscopy* 2005;21(4):385

References

34. Walker JM. Histological Study of the Fetal Development of the Human Acetabulum and Labrum: Significance in Congenital Hip Disease. *Yale J Bio Med* 1981;54(4):255.
35. Petersilge C et al. Acetabular Labral Tears: Evaluation with MR Arthrography. *Radiology* 1996;200:231-35.
36. Czerny C et al. MR Arthrography of Adult Acetabular Capsular-Labral Complex. *AJR* 1999;173:345-49.
37. Dinuer P et al. Sublabral Sulcus at the Posteroinferior Acetabulum: A Potential Pitfall in MR Arthrography Diagnosis of Acetabular Labral Tears. *AJR* 2004;183:1745-53.
38. Keene GS, Villar RN. Arthroscopic anatomy of the hip: an in vitro study. *Arthroscopy* 1994;10:392-399
39. Saddik D et al. Prevalence and Location of Acetabular Sublabral Sulci at Hip Arthroscopy with Retrospective MRI Review. *AJR* 2006;187:W507-511.
40. Studler U et al. MR Arthrography of the Hip: Differentiation between an Anterior Sublabral Recess as a Normal Variant and a Labral Tear. *Radiology* 2008;249(3):947
41. Byrd T. Hip Arthroscopy: The Supine Approach. *Tech in Ortho* 2005;20(1):17
42. Atlihan D et al. Arthroscopic Treatment of a Symptomatic Hip Plica. *Clin Ortho Rel Res* 2003;411:174-177.
43. Stoller D et al. *Magnetic Resonance Imaging in Orthopaedics and Sports Medicine* 3rd edition. Lippincott Williams & Wilkins 2007
44. Resnick D et al. *Internal Derangements of Joints* 2nd edition. Elsevier 2007.
45. Resnick D. IDJ Advanced Intensive MR Imaging Course Feb 2009.